

Ready to take the  
**NEXT  STEP**  
in treating your **ITP**?

These questions can help you and your doctor discuss your immune thrombocytopenia (ITP) treatment goals and whether WAYRILZ™ (rilzabrutinib) is right for you.

What is the **ideal platelet level** for me?

Is it possible to have **long-lasting control over my platelet levels**?

Can we discuss how ITP has been **impacting my quality of life** in ways such as \_\_\_\_\_?  
(fill in the blank).

Why may I still be **experiencing symptoms like fatigue** even when my platelets are in my target range?

How is **targeting key sources of ITP with WAYRILZ** different than other ITP treatments?

What **kind of side effects** have people **experienced** with WAYRILZ?

## Is WAYRILZ an option for me?

All images shown are patient portrayals.

### WHAT IS WAYRILZ?

WAYRILZ is a prescription medicine that is used to treat adults with persistent or chronic immune thrombocytopenia (ITP) who received a prior treatment that did not work well enough.

It is not known if WAYRILZ is safe and effective in children.

Please see Important Safety Information throughout and in full on page 3, and the full Prescribing Information, including Patient Information.

**WAYRILZ**<sup>™</sup>  
(rilzabrutinib) 400 mg  
tablets

## WAYRILZ:



### Improves and sustains platelet counts

By month 6 in the clinical study, **23% of people taking WAYRILZ reached and maintained target platelet counts** (at least 50,000) compared to 0% of people not taking WAYRILZ.

During the last 12 weeks of the study, people reached a platelet level of 50,000 or higher more than 5 times when measured weekly, without rescue therapy, including at least twice in the final 6 weeks.



### Designed to target key sources of ITP

WAYRILZ works in multiple ways to protect platelets and restore immune balance.



### Has an established safety profile in the clinical study

The most common side effects included diarrhea, nausea, headache, abdominal pain, and COVID-19.

## People taking WAYRILZ reported impact on quality of life

At month 6, in the same questionnaire taken at the start of the study (the ITP-PAQ), patients rated impact on different aspects of their lives. Each measurement was reported with a score ranging from 0 to 100, with a higher score indicating improving (▲) and lower score indicating worsening (▼).

	People taking WAYRILZ	People not taking WAYRILZ		People taking WAYRILZ	People not taking WAYRILZ
Overall Quality of Life	▲ +10.6	▲ +2.3	Fatigue/Sleep	▲ +9.3	▼ -5.0
Psychological Health	▲ +11.9	▼ -1.0	Fear	▲ +7.8	▼ -5.0
Bother-Physical Health	▲ +10.0	▼ -4.7	Symptoms	▲ +10.3	▲ +2.1
Social Activity	▲ +9.4	▲ +1.3	Women's Reproductive Health	▲ +5.8	▼ -16.7
Activity	▲ +9.96	▼ -1.3	Work	▲ +4.9	▲ +3.8

The results of these analyses are descriptive and were not statistically tested to determine a difference between treatment groups.

Share these pages with your doctor to decide if [WAYRILZ](#) may be right for you.

### IMPORTANT SAFETY INFORMATION

#### What should I tell my healthcare provider before taking WAYRILZ?

Tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems • have kidney problems
- are pregnant or plan to become pregnant. WAYRILZ may harm your unborn baby. If you are able to have a baby, your healthcare provider will do a pregnancy test before starting treatment with WAYRILZ. Females who are able to become pregnant should use an effective birth control during treatment with WAYRILZ and for 1 week after the last dose.

Please see Important Safety Information throughout and in full on page 3, and the full [Prescribing Information](#), including [Patient Information](#).

**WAYRILZ**<sup>TM</sup>  
(rilzabrutinib) 400 mg tablets

## WHAT IS WAYRILZ?

WAYRILZ is a prescription medicine that is used to treat adults with persistent or chronic immune thrombocytopenia (ITP) who received a prior treatment that did not work well enough.

It is not known if WAYRILZ is safe and effective in children.

## IMPORTANT SAFETY INFORMATION

### What should I tell my healthcare provider before taking WAYRILZ?

**Tell your healthcare provider about all of your medical conditions, including if you:**

- have liver problems
- have kidney problems
- are pregnant or plan to become pregnant. WAYRILZ may harm your unborn baby. If you are able to have a baby, your healthcare provider will do a pregnancy test before starting treatment with WAYRILZ. Females who are able to become pregnant should use an effective birth control during treatment with WAYRILZ and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with WAYRILZ and for at least 1 week after the last dose.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking WAYRILZ with certain other medicines may affect how WAYRILZ works and can cause side effects. WAYRILZ may also affect how other medicines work.

### What should I avoid while taking WAYRILZ?

You should avoid grapefruit, starfruit and products that have these fruits, and Seville oranges (often used in marmalades) during treatment with WAYRILZ. These products may increase the amount of WAYRILZ in your blood, which increases the risk of side effects of WAYRILZ.

### What are the possible side effects of WAYRILZ?

**WAYRILZ may cause serious side effects, including:**

- **Serious Infections.** WAYRILZ can increase the risk of infections, including serious infections that can lead to death. Your healthcare provider will check you for signs and symptoms of infection during your treatment with WAYRILZ. Tell your healthcare provider right away if you get any signs or symptoms of infection, including fever, chills, or flu-like symptoms.
- **Liver problems including Drug-Induced Liver Injury (DILI).** Liver problems, which may be severe, life-threatening, or lead to death have happened in people treated with Bruton's tyrosine kinase (BTK) inhibitors. Your healthcare provider will do blood tests to check your liver before and as necessary during treatment with WAYRILZ. Tell your healthcare provider right away if you have any signs or symptoms of liver problems, including stomach-area (abdominal) pain or discomfort, dark or "tea-colored" urine, or yellowing of the skin or the white part of your eyes.

**The most common side effects of WAYRILZ include** diarrhea, nausea, headache, stomach area (abdominal) pain, and COVID-19.

These are not all of the possible side effects of WAYRILZ.

**Please see full [Prescribing Information](#), including [Patient Information](#).**

**sanofi**

© 2025 Sanofi. All rights reserved.

WAYRILZ and Sanofi are trademarks of Sanofi or an affiliate.

MAT-US-2504565-v1.0-09/2025

**WAYRILZ**<sup>™</sup>  
(rilzabrutinib) 400 mg  
tablets