

The NEXT STEP

Forward in ITP Treatment



WAYRILZ[™]
(rilzabrutinib) 400 mg
tablets

FOR ADULTS with CHRONIC ITP

WAYRILZ offers long-lasting platelet control* and was studied for impact on patient-reported quality of life.

All images shown within are patient portrayals.

*During the last 12 weeks of the study, people reached a platelet level of 50,000 or higher more than 5 times when measured weekly, without rescue therapy, including at least twice in the final 6 weeks.

WHAT IS WAYRILZ?

WAYRILZ is a prescription medicine that is used to treat adults with persistent or chronic immune thrombocytopenia (ITP) who received a prior treatment that did not work well enough.

It is not known if WAYRILZ is safe and effective in children.

Please see Important Safety Information on page 14, and the full Prescribing Information, including Patient Information.

Platelet Counts Don't Capture the Full Reality of ITP

It starts with understanding what's happening inside your body

Immune thrombocytopenia (ITP) is a rare autoimmune disorder in which the immune system is out of balance in more ways than one. This causes your body to mistakenly attack its own platelets.

Platelets are vital blood cells that help your blood clot. When platelets are destroyed or reduced, it leads to a lower-than-normal platelet count, raising the risk of bleeding.

Bleeding symptoms of ITP include:



Easy bruising or bruises without a clear cause




Red or purple pinpoint spots on arms and legs (petechiae)



Frequent or prolonged bleeding (like nose bleeds, heavy periods, and bleeding gums)

ITP can also impact how you feel

Over 1500 people with ITP participated in a large global study. They reported:



INTRODUCING WAYRILZ

Take the Next Step Forward in Your ITP Treatment



Long-lasting platelet control

WAYRILZ improves and sustains platelet counts.

During the last 12 weeks of the study, 23% of people taking WAYRILZ (vs 0% of those who did not) reached a platelet level of 50,000 or higher more than 5 times when measured weekly, without rescue therapy, including at least twice in the final 6 weeks.



Measured impact to patient-reported quality of life

During the study, people evaluated impact to parts of their daily life, including fatigue, emotional health, and more.

Not a steroid. No infusions, no injections. Just one pill, twice a day.

IMPORTANT SAFETY INFORMATION

What should I tell my healthcare provider before taking WAYRILZ?

Tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have kidney problems

WAYRILZTM
(rilzabrutinib) 400mg tablets

Please see Important Safety Information on page 14, and the full [Prescribing Information](#), including [Patient Information](#).

WAYRILZ is designed to target key sources of ITP

With ITP, a protein called **BTK (Bruton's tyrosine kinase)** plays a key role in your immune system's attack on your platelets by sending signals that lead to platelet destruction.

WAYRILZ is a BTK inhibitor, meaning it blocks those harmful signals at the source, and works in multiple ways to protect platelets by:

 **Preventing** autoantibodies from marking platelets for destruction

 **Stopping** immune cells from attacking and breaking down platelets

**WAYRILZ helps restore immune balance,
which may allow your body to better protect
and raise platelet counts.**

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all of your medical conditions, including if you (cont'd):

- are pregnant or plan to become pregnant. WAYRILZ may harm your unborn baby. If you are able to have a baby, your healthcare provider will do a pregnancy test before starting treatment with WAYRILZ. Females who are able to become pregnant should use an effective birth control during treatment with WAYRILZ and for 1 week after the last dose.

How WAYRILZ Was Studied

The clinical study focused on adults with persistent or chronic ITP who didn't respond well to or couldn't tolerate previous treatments. It compared **133 people** who took WAYRILZ to **69 people** who did not for a total of 24 weeks (or 6 months).

What does it mean to be a "Responder"?



Responders were patients whose platelet counts improved by month 3 of the study. These patients continued to receive treatment for the full 6 month study.

This meant reaching a platelet response (at least 50,000 or between 30,000 and 50,000 and doubling from their starting level) without needing rescue treatments.

64%

of people taking WAYRILZ were responders compared to 32% of people not taking WAYRILZ.

IMPORTANT SAFETY INFORMATION


Tell your healthcare provider about all of your medical conditions, including if you (cont'd):

- are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with WAYRILZ and for at least 1 week after the last dose.

Long-lasting Platelet Control With WAYRILZ

BY MONTH 6:

WAYRILZ improved and sustained platelet counts.*



*During the last 12 weeks of the study, people reached a platelet level of 50,000 or higher more than 5 times when measured weekly, without rescue therapy, including at least twice in the final 6 weeks.

Platelet counts rose quickly

People taking WAYRILZ (133 people) **reached target platelet levels in 36 days.** Those not taking WAYRILZ (69 people) took longer or didn't reach target levels by the end of the study.†

Responders (85 people) **reached target platelet levels typically around 2 weeks.**

†During the study, these people reached a platelet level of 50,000 or higher or between 30,000 and less than 50,000 and at least doubled from starting level, without rescue therapy.

Platelets stayed up week after week with WAYRILZ

7 weeks with WAYRILZ

5 days without WAYRILZ

10X LONGER

By month 6, people taking WAYRILZ kept platelet counts at desired levels ($\geq 50,000$ or between 30,000 and 50,000 and doubled from baseline) for a longer time compared to those not taking WAYRILZ.

People taking WAYRILZ needed fewer rescue treatments

During the study, medicines like IVIg, or intravenous immunoglobulin, were given only if needed. **58% of people who did not take WAYRILZ needed additional treatment** compared to only 33% of people taking WAYRILZ.

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking WAYRILZ with certain other medicines may affect how WAYRILZ works and can cause side effects. WAYRILZ may also affect how other medicines work.

WAYRILZ™
(rilzabrutinib) 400 mg tablets

Please see Important Safety Information on page 14, and the full [Prescribing Information](#), including [Patient Information](#).

People Taking WAYRILZ Reported Impact to Quality of Life



The ITP-PAQ

People in the study completed a survey called the ITP-PAQ that looked at how ITP affected different parts of their lives.

IMPORTANT SAFETY INFORMATION

What should I avoid while taking WAYRILZ?

You should avoid grapefruit, starfruit and products that have these fruits, and Seville oranges (often used in marmalades) during treatment with WAYRILZ. These products may increase the amount of WAYRILZ in your blood, which increases the risk of side effects of WAYRILZ.



Please see Important Safety Information on page 14, and the full [Prescribing Information](#), including [Patient Information](#).

How impact to quality of life was evaluated

At month 6, in the same questionnaire taken at the start of the study (the ITP-PAQ), patients rated impact on different aspects of their lives. Each measurement was reported with a score ranging from 0 to 100, with a higher score indicating improving (∧) and lower score indicating worsening (∨).

	TAKING WAYRILZ	NOT TAKING WAYRILZ
Overall Quality of Life	∧ +10.6	∧ +2.3
Psychological Health	∧ +11.9	-1.0 ∨
Bother-Physical Health	∧ +10.0	-4.7 ∨
Social Activity	∧ +9.4	∧ +1.3
Activity	∧ +9.96	-1.3 ∨
Fatigue/Sleep	∧ +9.3	-5.0 ∨
Fear	∧ +7.8	-5.0 ∨
Symptoms	∧ +10.3	∧ +2.1
Women's Reproductive Health	∧ +5.8	-16.7 ∨
Work	∧ +4.9	∧ +3.8

The results of these analyses are descriptive and were not statistically tested to determine a difference between treatment groups.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of WAYRILZ?

WAYRILZ may cause serious side effects, including:

- **Serious infections.** WAYRILZ can increase the risk of infections, including serious infections that can lead to death. Your healthcare provider will check you for signs and symptoms of infection during your treatment with WAYRILZ. Tell your healthcare provider right away if you get any signs or symptoms of infection, including fever, chills, or flu-like symptoms.

WAYRILZ Has an Established Safety Profile in the Clinical Study

When starting a new treatment, it's natural to have questions, especially about safety. Here's a look at what some people have experienced during the clinical study.

The most common side effects of WAYRILZ include:

- diarrhea
- stomach area (abdominal) pain
- nausea
- COVID-19
- headache

If you experience diarrhea, nausea, or stomach area (abdominal pain) during treatment with WAYRILZ, taking it with food may reduce these side effects.

WAYRILZ may cause serious side effects, including serious infections.

Tell your healthcare provider right away if you get any signs or symptoms of infection, including fever, chills, or flu-like symptoms.

Severe liver problems, including Drug-Induced Liver Injury, have happened in people treated with Bruton's tyrosine kinase (BTK) inhibitors.

In ITP, some people in the WAYRILZ clinical trial had mild-to-moderate increases in liver enzyme levels. Liver enzymes are something doctors monitor during treatment to check how your liver is working. Your healthcare team will regularly check your liver function to help keep you safe and supported throughout your treatment.

These are not all of the possible side effects of WAYRILZ. Call your doctor for medical advice, including tips on managing potential side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Convenient Oral Dosing That Fits in Your Schedule



Take WAYRILZ *with* or *without* food
Swallow pills whole, with water. Do not chew, cut, or crush.

Take WAYRILZ exactly as your doctor tells you to take it. Do not change your dose or stop taking WAYRILZ unless your doctor tells you to.

Please tell your doctor if you are taking medications for heartburn, acid reflux, or ulcers, such as antacids, H2 (histamine-2) blockers, or proton pump inhibitors (PPIs), as these may require additional instructions.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of WAYRILZ?

- **Liver problems including Drug-Induced Liver Injury (DILI).** Liver problems, which may be severe, life-threatening, or lead to death have happened in people treated with Bruton's tyrosine kinase (BTK) inhibitors. Your healthcare provider will do blood tests to check your liver before and as necessary during treatment with WAYRILZ. Tell your healthcare provider right away if you have any signs or symptoms of liver problems, including stomach-area (abdominal) pain or discomfort, dark or "tea-colored" urine, or yellowing of the skin or the white part of your eyes.

Support That Stays With You

Sanofi Support for WAYRILZ

sanofi *Support* | HemAssist™

After your doctor prescribes WAYRILZ, **HemAssist Sanofi Support** can assist with access to WAYRILZ, financial assistance, and education throughout your treatment journey.

Support from HemAssist:



Financial assistance

Every person's situation is different, so we work directly with you to help identify the financial assistance programs for which you may be eligible.



Treatment education

We can help you understand more about treatment with WAYRILZ by connecting you with a dedicated Clinical Educator.



Access to treatment

Our team works with you and your care team to verify your insurance coverage and get you started on WAYRILZ as efficiently as possible.

Your HemAssist team

Clinical Educator (CE)

Your CE is your point of contact for questions and tips about WAYRILZ, taking and tracking your treatment, and ongoing support.

Case Manager (CM)

Your CM is available to assist with navigating insurance coverage and identifying available financial support for eligible patients.

Your team will not provide personal medical advice. For all questions about your individual health needs, please speak with your doctor.

Ready to enroll in HemAssist?

Visit SanofiHemAssist.com or give us a call at **1-833-723-5463 Monday – Friday 8 AM to 7 PM ET.**

IMPORTANT SAFETY INFORMATION

What are the possible side effects of WAYRILZ?

The most common side effects of WAYRILZ include diarrhea, nausea, headache, stomach area (abdominal) pain, and COVID-19.

These are not all of the possible side effects of WAYRILZ.



Please see Important Safety Information on page 14, and the full [Prescribing Information](#), including [Patient Information](#).



Ready to take the next step with WAYRILZ?

Receive a personalized guide to help support discussions with your doctor about the next steps in your treatment plan.

Answer a few easy, multiple-choice questions about your symptoms and treatment goals, and get results tailored to your needs.

START HERE and get your personalized guide!

WAYRILZ[™]
(rilzabrutinib) 400 mg
tablets

WHAT IS WAYRILZ?

WAYRILZ is a prescription medicine that is used to treat adults with persistent or chronic immune thrombocytopenia (ITP) who received a prior treatment that did not work well enough. It is not known if WAYRILZ is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What should I tell my healthcare provider before taking WAYRILZ?

Tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have kidney problems
- are pregnant or plan to become pregnant. WAYRILZ may harm your unborn baby. If you are able to have a baby, your healthcare provider will do a pregnancy test before starting treatment with WAYRILZ. Females who are able to become pregnant should use an effective birth control during treatment with WAYRILZ and for 1 week after the last dose.
- are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with WAYRILZ and for at least 1 week after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking WAYRILZ with certain other medicines may affect how WAYRILZ works and can cause side effects. WAYRILZ may also affect how other medicines work.

What should I avoid while taking WAYRILZ?

You should avoid grapefruit, starfruit and products that have these fruits, and Seville oranges (often used in marmalades) during treatment with WAYRILZ. These products may increase the amount of WAYRILZ in your blood, which increases the risk of side effects of WAYRILZ.

What are the possible side effects of WAYRILZ?

WAYRILZ may cause serious side effects, including:

- **Serious infections.** WAYRILZ can increase the risk of infections, including serious infections that can lead to death. Your healthcare provider will check you for signs and symptoms of infection during your treatment with WAYRILZ. Tell your healthcare provider right away if you get any signs or symptoms of infection, including fever, chills, or flu-like symptoms.
- **Liver problems including Drug-Induced Liver Injury (DILI).** Liver problems, which may be severe, life-threatening, or lead to death have happened in people treated with Bruton's tyrosine kinase (BTK) inhibitors. Your healthcare provider will do blood tests to check your liver before and as necessary during treatment with WAYRILZ. Tell your healthcare provider right away if you have any signs or symptoms of liver problems, including stomach-area (abdominal) pain or discomfort, dark or "tea-colored" urine, or yellowing of the skin or the white part of your eyes.

The most common side effects of WAYRILZ include diarrhea, nausea, headache, stomach area (abdominal) pain, and COVID-19.

These are not all of the possible side effects of WAYRILZ.



Please see the full [Prescribing Information](#), including [Patient Information](#).

Talk to your doctor about
**moving forward
with WAYRILZ**



All images shown within are patient portrayals.



Long-lasting platelet control

WAYRILZ improves and sustains platelet counts. During the last 12 weeks of the study, 23% of people taking WAYRILZ (vs 0% of those who did not) reached a platelet level of 50,000 or higher more than 5 times when measured weekly, without rescue therapy, including at least twice in the final 6 weeks.



Measured impact to patient-reported quality of life

During the study, people evaluated impact to parts of their daily life, including fatigue, emotional health, and more.



Designed to target key sources of ITP

WAYRILZ works in multiple ways to protect platelets and restore immune balance.

Not a steroid. No infusions, no injections. Just one pill, twice a day.

Explore more about **WAYRILZ** and [sign up today](#).

IMPORTANT SAFETY INFORMATION

What should I tell my healthcare provider before taking WAYRILZ?

Tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems
- have kidney problems
- are pregnant or plan to become pregnant. WAYRILZ may harm your unborn baby. If you are able to have a baby, your healthcare provider will do a pregnancy test before starting treatment with WAYRILZ. Females who are able to become pregnant should use an effective birth control during treatment with WAYRILZ and for 1 week after the last dose.

Please see Important Safety Information on page 14, and the full [Prescribing Information](#), including [Patient Information](#).

sanofi

© 2025 Sanofi. All rights reserved.
WAYRILZ, HemAssist, and Sanofi are trademarks of Sanofi or an affiliate.
MAT-US-2508676-v1.0-09/2025

WAYRILZ[™]
(rilzabrutinib) 400 mg
tablets